

PESTICIDE LABELS

TRIMEC 992

ECIMEN LABEL – SPECIMEN LABEL – SPECIMEN LABEL – SPECIMEN LABEL – SPECIMEN LABEL – SPEC

TRIMEC 992 BROADLEAF HERBICIDE

Controls Dandelion, Clover, Knotweed, Henbit,
Chickweed, Plantain, Spurge, and
Other Listed Broadleaf Weeds

Use this product only in accordance with its
labeling and with the Worker Protection Standard,
40 CFR Part 170.

ACTIVE INGREDIENTS:
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid 30.56%
Dimethylamine salt of (+)-R)-2-(2-methyl-4-chlorophenoxy)
propionic acid 8.17%
Dimethylamine salt of dicamba-3,5-dichloro-o-anisic acid 2.77%
OTHER INGREDIENTS 58.50%
TOTAL 100.00%

THIS PRODUCT CONTAINS:
2.38 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 25.38%
0.83 lb. (+)-R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent
per gallon or 6.75%
0.23 lb. 3,5-dichloro-o-anisic acid equivalent per gallon or 2.30%
Isomer Specific by ADAC Methods.

KEEP FROM FREEZING

KEEP OUT OF REACH OF CHILDREN DANGER - PELIGRO

Si Usted no entiende la etiqueta, busque a alguien para que se la
explique a Usted en detalle. (If you do not understand the label, find
someone to explain it to you in detail.)

STOP READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER: Corrosive. Causes irreversible eye damage. Harmful if
swallowed. Harmful if absorbed through skin. Do not get in eyes, on
skin, or on clothing. Wash thoroughly with soap and water after
handling and before eating, drinking, chewing gum, using tobacco or
using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are natural
rubber. If you want more options, follow the instructions for category A
on an EPA chemical-resistance category selection chart.

All mixes, loaders, applicators and other handlers must wear*:

- protective eyewear,
- long-sleeved shirt and long pants,
- shoes and socks,
- chemical-resistant gloves and
- chemical-resistant apron when mixing or loading, cleaning up spills
or equipment, or otherwise exposed to the concentrate.

*Applications may choose not to wear protective eyewear OSHA with dilution rates
greater (higher) than 5:1 or greater (higher) than 5 parts of water to 1 part
of product.

See engineering controls for additional requirements.

Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that
meets the requirements listed in the Worker Protection Standard
(WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the
handler PPE requirements may be reduced or modified as specified in
the WPS.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no
such instructions for washables exist, use detergent and hot water.
Keep and wash PPE separately from other laundry. Discard clothing
and other absorbent material that have been drenched or heavily
contaminated with the product's concentrate. Do not reuse them.

User Safety Recommendations

*Users should wash hands before eating, drinking, chewing gum,
using tobacco, or using the toilet.
*Users should remove clothing/PPE immediately if pesticide gets
inside. Then wash thoroughly and put on clean clothing. If pesticide
gets on skin, wash immediately with soap and water.
*Users should remove PPE immediately after handling this product.
Wash the outside of gloves before removing. As soon as possible,
wash thoroughly and change into clean clothing.

First Aid

If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or on clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison
control center or doctor or going for treatment. You may also contact
1-877-800-5556 for emergency medical treatment advice.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate
the use of gastric lavage.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may
adversely affect non-target plants. Do not apply directly to water, to
areas where surface water is present, or to intertidal areas below the
mean high water mark. Drift and runoff may be hazardous to aquatic
organisms in water adjacent to treated areas. Do not contaminate
water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with
chemicals detected in groundwater. The use of this chemical in areas
where soils are permeable, particularly where the water table is
shallow, may result in groundwater contamination. Application around a
cistern or well may result in contamination of drinking water or
groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsis-
tent with its labeling.

Do not apply this product in a way that will contact workers or other
persons, either directly or through drift. Only protected handlers may be
in the area during application. For any requirements specific to your
State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the
Worker Protection Standard, 40 CFR part 170.

This standard contains requirements for the protection of agricultural
workers on farms, forests, nurseries, and greenhouses, and handlers of
agricultural pesticides. It contains requirements for training, decon-
tamination, notification, and emergency assistance. It also contains
specific instructions and exceptions pertaining to the statements or

(cont. on next page)

[LINK TO LABEL](#)

24-0-5 PLANT NUTRIENT

Plant Nutrient 24-0-5

GUARANTEED ANALYSIS

Total Nitrogen (N)	24.00%
24.00% Urea Nitrogen*	5.00%
Soluble Potash (K ₂ O).....	4.00%
Iron (Fe).....	

Derived From: Iron Sulfate, Muriate of Potash, Polymer coated urea, and Urea.

Salt Index: 3042 Bulk Density: 63.81

Net Weight 50 lbs (22.70 kg)

50 lbs. Covers 12,000 sq. ft.

(These rates equal 1 lb. nitrogen per 1,000 sq. ft.)

*The Urea in this product has been coated to provide 12.00% Slow-Release Nitrogen.

ALSO CONTAINS NON-PLANT FOOD INGREDIENTS

Paenibacillus polymyxa 400,000 CFU per gram,
Bacillus subtilis 300,000,000 cfu per gram,
Bacillus pumilus 300,000 cfu per gram,
Bacillus licheniformis 300,000 cfu per gram,
Bacillus amyloliquefaciens 300,000 cfu per gram,
Bacillus megaterium 300,000 cfu per gram,
Pseudomonas putida 200,000 cfu per gram,
Pseudomonas fluorescens 200,000 cfu per gram,
Pseudomonas aureofaciens 200,000 cfu per gram,
Azospirillum amazonense 200,000 cfu per gram,
Azospirillum lipoferum 200,000 cfu per gram
Expiration date: 1-1-2020

Item Number: 2034623T

NUTRIENT CALCULATIONS WITH SPREADER SETTINGS

Product	Low Rate	Medium Rate	High Rate
	2.08 lbs/M	3.12 lbs/M	4.16 lbs/M
24-0-5			
Lbs Nutrients/M	N: 50	N: 75	N: 100
	P: 00	P: 00	P: 00
	K: 10	K: 15	K: 20
Rotary Spreader	Settings	Settings	Settings
	3	4	5
Accugen		K	M
Earthway 2400	12	13	15
Lesco 00600	14	16	18
Proclaim	H	I	K
Spyker	4	4.5	5

PA17240115535

Powered by Sanctuary Detonation

Directions – Please Read
Carefully Before Using
General Information

This product is a highly versatile premium mini-sized fertilizer that is ideal for the nutrient demands of top quality turf management. The precise balance of nutrients promotes a superior, sustained color response that produces the visual appeal turf professionals desire.

How to Use*

For optimum performance, begin spring applications just before soil temperatures reach 50 degrees F. Although nitrogen can be applied at the rate of 1.0 lb Per 1,000 sq ft every 8 weeks, it is also recommended that lighter, half-rate applications be made each month, especially during hot, dry periods. Continue scheduled feedings throughout the growing season until soil temperature retreats to below 50 degrees F. Late fall and dormant feedings are recommended after soil temperatures remain below 50 degrees F for several consecutive days and turf growth has slowed considerably.

Apply product evenly with a calibrated spreader. For best results, irrigating after applications will aid in granule penetration in close-cut dense turf. Removing grass catchers for initial mowing will minimize particle pick-up.

*Southern turf grasses, including St. Augustine and hybrid Bermuda grasses may require higher levels of nitrogen for optimum performance.

NEVER FERTILIZE WHEN THE GRASS IS MOIST
ALWAYS CLEAN OUT SPREADER AFTER EACH USE.

Note: These Settings are approximate. Due to atmospheric conditions and wear on spreader it is impossible to be 100% accurate. It is wise to set your spreader to the rate and then measure the coverage of only one pound of test feed. If you are under applying, increase the rate. If you are over applying, decrease the rate. If you think it's better to under apply than to over apply use formulas with more controlled releases, especially during hot dry conditions.

Note: These Settings are Approximate and can be influenced by other contributing factors Vogel Seed & Fertilizer Inc. makes no warranty as to the uniformity of coverage actually obtained from the settings listed.

KEEP OUT OF REACH OF CHILDREN

Caution: May cause irritation. On Contact with skin or eyes, flush with plenty of water.

DISTRIBUTED BY VOGEL SEED & FERTILIZER INC.
1891 SPRING VALLEY ROAD, JACKSON, WI 53037

20-0-0 + SOLU CAL

Solu-Cal^{USA}

20-0-0 + SOLU-CAL

With Slow Release XCU Nitrogen
Coarse Grade (SGN 210)

20-0-0 Guaranteed Analysis

Total Nitrogen (N).....20.0%

20.0% Urea Nitrogen*

Calcium (Ca).....21.0%

Derived from: Urea, Polymer-coated sulfur-coated urea, Calcium Carbonate

*6.0% slowly available urea nitrogen from polymer-coated, sulfur-coated urea

ALSO CONTAINS NON-PLANT FOOD INGREDIENT:

Soil Amending Guarantee Analysis:

Active Ingredients:

8.00% Lignin Sulfonate.

0.50% Microbes

Baenobacillus polymyxa.....150,000 cfu per gram
Bacillus subtilis.....150,000 cfu per gram
Bacillus pumilus.....150,000 cfu per gram
Bacillus sphaeriformis.....150,000 cfu per gram
Bacillus amyloliquefaciens.....150,000 cfu per gram
Bacillus megaterium.....150,000 cfu per gram
Asospirillum brassicicola.....150,000 cfu per gram
Asospirillum lipoferrum.....150,000 cfu per gram
Trichoderma harzianum.....150,000 cfu per gram

Inert Ingredients (as a soil amendment):

91.50% Fertilizer Materials

Purpose: microbes decompose organic materials in the soil.

Information concerning the raw materials composing this product can be obtained by writing to the Guarantor listed below, and referring the batch number found on this container. Information regarding the contents & levels of metals in this product is on the internet at <http://www.aagfco.com/metals.htm>

Net Wt. 50 lbs. (22.68 kg)

Directions For Use. Apply only as directed.

SOLU-CAL 20-0-0 may be applied to a wide variety of turf grasses & newly seeded turf areas. For best results water in after Application.

- **FOR EXISTING TURF:** Coverage 10 000 sq ft. Apply 5 lbs per 1000 sq ft or 220 lbs per acre. To assure complete and even coverage apply at half rates in two directions. Apply to dry turf. Water in thoroughly.

SOLU-CAL 20-0-0 if used at 220 lbs per acre will provide 1 lb of nitrogen and 1.0 lb of calcium per 1000 sq ft with each application. It is ideal for all turf types.

Store in a cool, dry place, do not contaminate water by storage or disposal.

EARTHWAY		CYCLONE		SCOTT'S A2000/A1000		LESCO		PRIZELAWN RE-CUR	
Set	width	Set	width	Set	width	Set	width	Set	Width
15	12	5.5	14	L		I	14	J	14

Settings are for 5.0 lbs rate. These spreader settings are approximate. Age, condition of spreader, speed, and contour of terrain may cause settings to vary. Always calibrate your spreader before use.

BUYER ASSUMES ALL RESPONSIBILITY FOR SAFETY AND USE NOT IN ACCORDANCE WITH DIRECTIONS. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THE PRODUCT. PLANT INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT. EVIDENCE OF SUCH FACTORS AT WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND SELLER'S CONTROL. IN NO CASE SHALL THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

FLORIDA APPLICATIONS: For use on golf courses or other athletic turf in FL, we recommend you follow SL191, "Recommendations for N, P, K & Mg for Golf Course and Athletic Field Fertilization Based on Metlich I Extractor", available on the internet at <http://ada.fda.usd.edu/SL191>.

For commercial application to urban turf or lawns in FL, we recommend you follow the Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries, available on the internet at <http://www.fgic.com/FloridaGreenIndustries/BestManagementPractices020602.pdf>.

Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden.



Warning

Caution: Irritation. Causes serious eye damage. May cause respiratory irritation. May cause damage to organs through prolonged or repeated exposure if inhaled. Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Do not breathe dust. Wash hands thoroughly after handling. Do not eat, drink, or smoke when using this product. Use only in well-ventilated areas. Wear protective gloves/goggles/eye protection when protection is required. Remove person to fresh air and keep comfortable for breathing. If in eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If inhaled: Get medical attention if you feel unwell. If on skin: Wash with plenty of water. Take off contaminated clothing and wash it before reuse. Get medical advice/attention if you feel unwell. Store locked up. Dispose of contents/container according to local regulations. See Safety Data Sheet for further details regarding safe use of product.

Made in USA

Distributed by: GROWMARK Inc., 1701 Towanda Avenue, Bloomington, IL 61701

[LINK TO LABEL](#)

TURFLON

TRICLOPYR GROUP 4 HERBICIDE



Turflon® Ester Ultra

ACCEPTED
VIA NOTIFICATION
LABEL NOT REVIEWED

SEPT 25 2020

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration

HERBICIDE

Doc ID: 569584

For the control of annual and perennial broadleaf weeds and kikuyugrass in ornamental turf, rangeland and permanent grass pastures, and non-crop areas including industrial manufacturing and storage sites, rights-of-way including electrical power lines, communication lines, pipelines, roadsides, railroads, fence rows, non-irrigation ditch banks, and in the establishment and maintenance of wildlife openings. Use on these sites may include application to grazed areas.

Active Ingredient:

triclopyr: 3,5,6-trichloro-2-pyridinylloxyacetic acid, butoxyethyl ester.....	60.45%
Other Ingredients.....	39.55%
Total.....	100.00%

Acid Equivalent: triclopyr – 43.46% - 4 lb/gal

Keep Out of Reach of Children CAUTION

Refer to inside of label booklet for additional information including Directions for Use.

NET CONTENTS 2.5 GAL

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-566 EPA Est. 464-MI-1-00323100 1905

®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268



Turflon® Ester Ultra

HERBICIDE

For the control of annual and perennial broadleaf weeds and kikuyugrass in ornamental turf, rangeland and permanent grass pastures, and non-crop areas including industrial manufacturing and storage sites, rights-of-way including electrical power lines, communication lines, pipelines, roadsides, railroads, fence rows, non-irrigation ditch banks, and in the establishment and maintenance of wildlife openings. Use on these sites may include application to grazed areas.

Active Ingredient:

triclopyr: 3,5,6-trichloro-2-pyridinylloxyacetic acid, butoxyethyl ester.....	60.45%
Other Ingredients.....	39.55%
Total.....	100.00%

Acid Equivalent: triclopyr – 43.46% - 4 lb/gal

Keep Out of Reach of Children CAUTION

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional information including Directions for Use.

TRICLOPYR GROUP 4 HERBICIDE

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-566

EPA Est. 464-MI-1-00323100 1905

[LINK TO LABEL](#)

®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

NET CONTENTS 2.5 GAL

SOLU CAL HUMIC +PLUS

IDEAL FOR: turf, landscapes,
trees & shrubs, and agricultural crops

Solu-Cal Humic +PLUS

The Power of Humic Acid
teamed with *Solu-Cal* Soluble Calcium
in a dust-free pellet

Powered by **INTENSIFY**
Soil Microbial BioCatalyst™

- Improves nutrient availability
- Provides essential soluble Calcium
- Delivers 11lbs. Humic Acid per acre

50lb bag covers 11,000 sq. ft.



NET WT. 50LB (22.68 kg)



Solu-Cal Humic +PLUS

Humic PLUS is a unique unique formulation of soluble Calcium and Humic Acids complexed to supply essential soluble Calcium and Carbon from highly oxidized lignite coal to the soil. Humic PLUS contains Humic Acid extracted by wet chemistry alkaline extraction to refine the natural benefits of organic plant matter humus. Impregnated with INTENSIFY Soil Microbial BioCatalyst to stimulate beneficial microbial activity in the soil. Humic PLUS will support increased soil microbial populations, which have been shown to reduce nutrient requirements.

- Increases plant uptake of nutrients
- Increases root mass surface area
- Delivers 11 lbs. Humic Acid per acre
- Increases seedling density

Store in original container in cool, dry place.

Apply to dry turf. Water in thoroughly after application.

APPLICATION RATES

Large Older Trees: 5 cups/caliper inch broadcast to drip line

Container Transplanting

1/2 gal. 1 tablespoon

1 gallon pot: 4 tablespoons

4 gallon pot: 4 tablespoons

Agriculture/Broadcast

200 lbs./acre (applies 12 lbs./acre rate of Humic Acid)

Soil rate: 25 lbs./acre/yr

Extinguish Turf, Germination & Seed Starting
coverage 13M or 250 lbs. acre. Recommendation is two
applications per year. Ideal timing is early spring and
early fall. (applies 11 lb./acre rate of Humic Acid)
Hedge Banding: coverage 2.5M or 200 lbs./acre
Ornamental, annual, perennial & rose planting beds
Apply 10 to 15,000 sq. ft.
Shrub: 2 cups broadcast around root zone

Guaranteed Analysis

Calcium (Ca).....39%

Derived from Calcium Carbonate.

ALSO CONTAINS NON-PLANT FOOD INGREDIENTS

Soil Amending/Soil Amendment:

Acetic acid……………100.00 lb. per gram

Biochar……………100.00 lb. per gram

Calcium……………100.00 lb. per gram

Carbon……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Humic……………100.00 lb. per gram

Spreader Settings

SHROUDCAST SPREADER

6.1 LBS. PER 1,000 SQ.FT.

BAG COVERS 11,000 SQ.FT.

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

5

Solu-Cal

FL Company License No: F1792

Guaranteed by SOLUCAL-USA LLC

P.O. BOX 278, WEST WASHINGTON, WA 02176

774-678-0288 - www.solu-cal.com

Made in USA



[LINK TO LABEL](#)

SOLU-CAL

IDEAL FOR: turf, landscapes, trees & shrubs, and wildlife food plots

Solu-Cal PLUS

Enhanced Calcitic Lime
Cal Calcitico Mejorado

Solu-Cal PLUS Enhanced Calcitic Lime



- Raises soil pH efficiently
- Combats acidic soils
- University tested & industry proven
- Saves time & money
- Low dust formula

Covers up to 10,000 sq. ft.

Powered by **INTENSIFY**
Soil Microbial BioCatalyst™

NET WT. 50LB (22.68 kg)




Solu-Cal PLUS Enhanced Calcitic Lime

Covers up to 10,000 sq. ft.

Solu-Cal PLUS

Enhanced Calcitic Lime
Cal Calcitico Mejorado

Application Rates

Apply product at a rate of 10 lbs per 1,000 sq. ft. in the absence of soil test results. For soil test results, see below.

10 to 15 lbs per 1,000 sq. ft.
Apply 10 to 15 lbs per 1,000 sq. ft. in the absence of soil test results. For soil test results, see below.

15 to 20 lbs per 1,000 sq. ft.
Apply 15 to 20 lbs per 1,000 sq. ft. in the absence of soil test results. For soil test results, see below.

20 to 25 lbs per 1,000 sq. ft.
Apply 20 to 25 lbs per 1,000 sq. ft. in the absence of soil test results. For soil test results, see below.

Recomendaciones en la Aplicación

Aplicar el producto a una tasa de 10 libras por 1,000 metros cuadrados en ausencia de resultados de análisis de suelo.

10 a 15 libras por 1,000 metros cuadrados.
Aplicar 10 a 15 libras por 1,000 metros cuadrados en ausencia de resultados de análisis de suelo.

15 a 20 libras por 1,000 metros cuadrados.
Aplicar 15 a 20 libras por 1,000 metros cuadrados en ausencia de resultados de análisis de suelo.

20 a 25 libras por 1,000 metros cuadrados.
Aplicar 20 a 25 libras por 1,000 metros cuadrados en ausencia de resultados de análisis de suelo.

For incorporation - after application wet thoroughly 30 minutes before tilling.

Garantizada Analisis

Weight (Net)	50.00 lbs
Net Weight (kg)	22.68 kg
Gross Weight (Net)	53.00 lbs
Gross Weight (kg)	24.04 kg
Net Weight (Net)	50.00 lbs
Net Weight (kg)	22.68 kg
Net Weight (Net)	50.00 lbs
Net Weight (kg)	22.68 kg
Net Weight (Net)	50.00 lbs
Net Weight (kg)	22.68 kg
Net Weight (Net)	50.00 lbs
Net Weight (kg)	22.68 kg


Fineness before Tilling

100 Mesh	100%
80 Mesh	100%
60 Mesh	100%
40 Mesh	100%
20 Mesh	100%

Spreaders Setting Chart Cálculo para programación de distribución


SPREADER SPREADER	WHEELS (in)	WHEELS (cm)
1000000	100	254
1000000	100	254
1000000	100	254
1000000	100	254
1000000	100	254
1000000	100	254
1000000	100	254
1000000	100	254
1000000	100	254

NET WT. 50LB (22.68 kg)



[LINK TO LABEL](#)

SANCTUARY 7-0-3



the Sanctuary
Working With Nature

SANCTUARY TDN[®] 7-0-3 TURF & ORNAMENTAL FERTILIZER

SANCTUARY TDN[®] 7-0-3 TURF & ORNAMENTAL FERTILIZER is a blend of protein meals. This product provides a slow release of nutrients that offers outstanding residual color, even growth, & turf density. This product is also an excellent pre-play heat nutrient application for dormant, ground cover and ornamental beds.

GUARANTEED ANALYSIS

Total Nitrogen (N).....	7.0%
0.2% Water Soluble Nitrogen	
6.8% Water Insoluble Nitrogen	
Soluble Potash (K ₂ O).....	3.0%
Calcium (Ca).....	3.0%

Derived from Feather Meal, Soybean Meal, Wheat Middings, Potassium Sulfate and Calcium Carbonate.

DIRECTIONS FOR USE

FOR GENERAL TURF MAINTENANCE PROGRAM:
Apply 10 lbs. per 1000 sq. ft. at a 12-week interval to optimize color, growth and rooting. During periods of stress apply 5 lbs. per 1000 sq. ft. at a 6-week interval.


FOR GENERAL ORNAMENTAL MAINTENANCE PROGRAM: Apply 5 lbs. per 100 sq. ft. in the spring and fall.

SPREADER SETTINGS

Spreader settings are an initial guideline for calibrating the spreader.

# RATE	LBS./1000 SQFT	LEAF	YARD	INCHES
1.00	14	8	54	11
0.50	7	8	40	5.5

TDN VALUE (TOTAL DIGESTIBLE NUTRIENT VALUE)



TDN Value is an important energy measure for the animal nutritionist and the producer. This energy is related to the value of nutrients which contribute to feed value. The TDN energy value is a relative measure of energy used in the product. The higher the TDN value, the more energy that is in the product. This released energy is the amount of energy available from the animal nutrition product. For example, all commercial feedstuffs have a TDN value and 100% moisture products have a minimum of 80 TDN value.

F 1386

Information regarding the contents and level of metals in this product is available on the internet at <http://www.aqfba.org/metal.htm>

THE SANCTUARY DIFFERENCE - TOTAL DIGESTIBLE NUTRIENT VALUE[®]

WARRANTY

The Sanctuary, Inc. warrants that this product conforms to the analysis on its label. We have used an accurate method with label directions, under normal conditions, this product is warranted for its intended purposes. Since timing, method of application, weather, plant, and soil conditions, interaction with other fertilizers, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

For additional information contact: Sanctuary210@rlypotash.com

Distributed by The Sanctuary
PO Box 656 • Winter Park, CO 80482
WWW.SANCTUARYPRODUCTS.COM

Net Weight:
40 LB. (18.2 KG)

SAN TDN 703-NY Label
REV 4/17

PRE-M® 0.86%

•PRE-M® 0.86% Plus Fertilizer

APPLICATION RATES, RATE AND APPLICATION CHART		SUGGESTED SPREADER SETTINGS	
Turfgrass Species	Weeds	Rate of 0.86% PRE-M Plus Fertilizer	Comments
SOUTHERN GRASSES Bermudagrass Bahiagrass Zoysiagrass Centipedegrass Tall Fescue	crabgrass broadleaf Poa annua barnyardgrass fall panicum wool prostrate spurge parietaria evening primrose big-leaf giantseeds	4.0 to 5.3 lb/1,000 sq ft 8.0 (14" x 20" broadcast) in the spring prior to weed germination 4.0 to 5.3 lb/1,000 sq ft 8.0 (14" x 20" broadcast) in the spring prior to weed germination	Apply a repeat application of 2.7 to 4.0 lb/1,000 sq ft in 18 - 274 foliar after a minimum of 4 weeks if extended control or where heavy weed infestations are expected.
NORTHERN GRASSES Kentucky Bluegrass Perennial Ryegrass Tall Fescue	crabgrass broadleaf Poa annua barnyardgrass fall panicum wool prostrate spurge parietaria evening primrose big-leaf giantseeds	4.0 to 5.3 lb/1,000 sq ft 8.0 (14" x 20" broadcast) in the spring prior to weed germination	Apply a repeat application of 2.7 to 4.0 lb/1,000 sq ft in 18 - 274 foliar after a minimum of 4 weeks if extended control or where heavy weed infestations are expected.
SOUTHERN GRASSES Bermudagrass Bahiagrass Zoysiagrass Centipedegrass Tall Fescue	crabgrass broadleaf Poa annua barnyardgrass fall panicum wool prostrate spurge parietaria evening primrose big-leaf giantseeds	4.0 to 5.3 lb/1,000 sq ft 8.0 (14" x 20" broadcast) in the spring prior to weed germination	Apply a repeat application of 2.7 to 4.0 lb/1,000 sq ft in 18 - 274 foliar after a minimum of 4 weeks if extended control or where heavy weed infestations are expected.
NORTHERN GRASSES Kentucky Bluegrass Perennial Ryegrass Tall Fescue	crabgrass broadleaf Poa annua barnyardgrass fall panicum wool prostrate spurge parietaria evening primrose big-leaf giantseeds	4.0 to 5.3 lb/1,000 sq ft 8.0 (14" x 20" broadcast) in the spring prior to weed germination	Apply a repeat application of 2.7 to 4.0 lb/1,000 sq ft in 18 - 274 foliar after a minimum of 4 weeks if extended control or where heavy weed infestations are expected.

*Not for use on greens or tees.
*Rates do not apply to Bermudagrass greens and tees.
DO NOT apply more than 5.3 lb of this product per 1,000 sq ft per application, and no more than 15.5 lb of this product per 1,000 sq ft per year.
The efficacy of this product will be improved if the application is followed by one half inch of water or its equivalent in sprinkler irrigation. If the product is not activated by rainfall or irrigation within 14 days, atritic weed control may result.

EPA REG. NO. 1040-1-00
EPA EST. NO. 82797-FL-001 (M)
First label of lot code indicates manufacturing site.
#088581

1560
Manufactured for: LESCO, Inc.
1385 East 36th Street
Cleveland, OH 44114-1114

Net Weight: 50 lb (22.7 kg)
(01) 00758073858119

•PRE-M® 0.86% Plus Fertilizer

Not for use on turf being grown for sale or for other commercial use as sod, or for commercial seed production, or for research purposes. For control of specified weeds in or on residential lawns, corporate lawns, parks, recreational areas, golf courses, cemeteries, and athletic fields.

- Prevents germination of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge, knotweed, and *Poa annua*.
- Prevents annual grassy and broadleaf weeds.

Active Ingredient:
Pendimethalin, N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzamine 0.86%
Other Ingredients: 99.14%
Total 100.00%
*Licensed under one or more of the following U.S. Patents: 3,920,742; 4,066,441.

0-0-7

MINI FERTILIZER

GUARANTEED ANALYSIS

SOLUBLE POTASH (K₂O) 7.00%
DERIVED FROM: Muriate of Potash.
CHLORINE (Cl) Max 5.25%

PRECAUTIONS
Use only on well-established turfgrasses with a dense and uniform stand. On newly planted areas, application should not be made until the turfgrass has filled in and been mowed at least four times. Do not use on turf that has been thinned or damaged due to winter injury, excessive moisture, etc., allow for turf recovery prior to making an application. Applications made to overseeded warm-season turfgrasses may cause thinning or injury of the overseeded species. Do not use on Bermudagrass or *Poa annua* (greens and tees) as injury may occur before reseeding or water overseeding treated turfgrasses for at least 2 months following application. Do not spray turfgrasses for 3 months after application.

CAUTION: Causes eye irritation. Harmful if absorbed through skin. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Avoid breathing dust.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

If swallowed	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by your poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact Chem-Trec 1-800-424-9300 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS
This product is toxic to fish. DO NOT apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. DO NOT contaminate water by cleaning of equipment or disposal of equipment washwaters or rinsate.

PERSONAL PROTECTIVE EQUIPMENT (PPE)
Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical resistant gloves made of any waterproof materials, such as butyl rubber > 14 mils, nitrile rubber > 14 mils, or neoprene rubber > 14 mils, shoes plus socks. Follow manufacturers' instructions for changing/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations
Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove PPE immediately after handling the product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Remove contaminated clothing and wash separately before reuse.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at time of pesticide application.
- Observe all cautions and limitations in this label and on the labels of products used in combination with this product. Do not use this product other than in accordance with the instructions set forth on this label. The use of this product not consistent with this label can result in injury to turf, animals, or persons. Keep container closed to avoid spills and contamination.
- DO NOT apply this product through any type of irrigation system.
- Do not enter or allow others to enter the treated area until dews have settled.
- Do not apply this product in a way that will contact any person or pets, either directly or by drift.
- Do not allow dust to drift from the application site and contact people, structures, people occupy at the time and the associated property, parks and recreation areas, nonlawn crops, aquatic and wetland areas, woodlands, pastures, rangelands, and animals.

GENERAL INFORMATION
This product is a combination pre-emergent herbicide and fertilizer formulated to provide control of most annual grasses and certain broadleaf weeds found in turfgrasses. This product is labeled for use on the following turfgrasses species: Kentucky Bluegrass, Perennial Ryegrass, Fine Fescue, Tall Fescue, Bermudagrass, Poa annua (except putting greens and tees), Bermudagrass, St. Augustinegrass, Bahiagrass, Centipedegrass, and Zoysiagrass.

APPLICATION
Apply product uniformly at the recommended rate using a properly calibrated spreader (see table below for approximate settings). The efficacy of this product will be improved if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation. This will remove the product from the turfgrass foliage and establish a pre-emergent herbicide barrier in the soil. Avoid contact of product with porous surfaces such as driveways, stone, or wood. Removing particles will help prevent staining.

PRODUCT CONTROLS
This product controls weeds as they germinate but will not control established weeds. Unusually cold, excessively wet, or hot and dry conditions that delay germination or extend germination over a long period of time can reduce weed control.

Applied according to label directions and under normal growing conditions, this product will not cause turf injury. Over-application can result in turf stand loss, turf injury, or soil residues. Uneven application can decrease weed control or cause turf injury.

Seedling diseases, cold weather, excessive moisture, high soil pH, high salt concentration, or drought can weaken turfgrasses and increase the possibility of turf damage from this product.

PROPER USE
Apply product uniformly at the recommended rate using a properly calibrated spreader (see table below for approximate settings). The efficacy of this product will be improved if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation. This will remove the product from the turfgrass foliage and establish a pre-emergent herbicide barrier in the soil. Avoid contact of product with porous surfaces such as driveways, stone, or wood. Removing particles will help prevent staining.

#088581
T

[LINK TO LABEL](#)

THREE-WAY ESTER II

38903 Lesco 3-Way (228-317-10404)(1g) LF 4/18/11 10:41 AM Page 2 (3,1)

LESCO

THREE-WAY™ ESTER II SELECTIVE HERBICIDE

A THREE-WAY POST-EMERGENT SELECTIVE BROADLEAF HERBICIDE FOR USE ON GOLF COURSES, PARKS, ORNAMENTAL TURF LAWNS, ROADSIDES, AND SIMILAR TURF AREAS.

CONTROLS (OR KILLS) DANDELIONS, CHICKWEEDS, PLANTAINS, OXALIS, SPURGE AND MANY OTHER BROADLEAF WEEDS, SOME OF WHICH ARE LISTED ON THIS LABEL.

CONTAINS MCPA, TRICLOPYR AND DICAMBA ONE GALLON COVERS UP TO 3.2 ACRES

ACTIVE INGREDIENTS:	
Isooctyl (2-ethylhexyl) Ester of 2-Methyl-4-Chlorophenoxyacetic Acid*	56.14%
Butoxyethanol Ester of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid**	5.00%
Dicamba (3,6-Dichloro-o-Anisic Acid)***	3.60%
OTHER INGREDIENTS	35.26%
TOTAL	100.00%
Isomer Specific AOAC Method, Equivalent to:	
*2-Methyl-4-Chlorophenoxyacetic Acid	36.0%, 3.0 lbs./gal.
**3,5,6-Trichloro-2-Pyridinyloxyacetic Acid	3.6%, 0.3 lbs./gal.
***3,6-Dichloro-o-Anisic Acid	3.6%, 0.3 lbs./gal.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL
PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

#069069

NET CONTENTS: 1 gal (3.78 L)

TM

EPA REG. NO. 228-317-10404

EPA EST. NO. 228-IL-1

Manufactured for: LESCO, Inc. • 1385 East 36th Street • Cleveland, OH 44114-4114

Job #38903

LESCO

THREE-WAY™ ESTER II SELECTIVE HERBICIDE

ACTIVE INGREDIENTS:	
Isooctyl (2-ethylhexyl) Ester of 2-Methyl-4-Chlorophenoxyacetic Acid*	56.14%
Butoxyethanol Ester of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid**	5.00%
Dicamba (3,6-Dichloro-o-Anisic Acid)***	3.60%
OTHER INGREDIENTS	35.26%
TOTAL	100.00%
Isomer Specific AOAC Method, Equivalent to:	
*2-Methyl-4-Chlorophenoxyacetic Acid	36.0%, 3.0 lbs./gal.
**3,5,6-Trichloro-2-Pyridinyloxyacetic Acid	3.6%, 0.3 lbs./gal.
***3,6-Dichloro-o-Anisic Acid	3.6%, 0.3 lbs./gal.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300.
For Medical Emergencies Only, Call (877) 325-1840.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none">• Immediately call a poison control center or doctor.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give any liquid to the person.• Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

NOTE TO PHYSICIAN
Contains Petroleum Distillates. May pose an aspiration pneumonia hazard.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always store pesticides in a secured warehouse or storage building. Containers should be opened in well-ventilated area. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable Containers 5 Gallons or Less: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

#069069

NET CONTENTS: 1 gal. (3.78 L)

EPA Reg. No. 228-317-10404

EPA Est. No. 228-IL-1

Manufactured for: LESCO, Inc.
1385 East 36th Street, Cleveland, OH 44114-4114

PEEL HERE TO OPEN

12726000

[LINK TO LABEL](#)

(R000100)

PF-21701-3

MERIT 75WSP



ACTIVE INGREDIENT:
*Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinone 75%
OTHER INGREDIENTS: 25%
100%

EPA Reg. No. 432-1318 EPA Est. No. Indicated by the 2nd and 3rd digits of the batch number on this package. (09)-53967-NJ-1 (05)-065387-AR-002
Keep water soluble packets in this container and store in a cool dry place but not below freezing (32° F).
Do Not Remove Packets From Container Except For Immediate Use

STOP - Read the label before use.
Keep out of reach of children.

CAUTION

PRECAUCION AL USUARIO: Si usted no puede leer o entender inglés, no use este producto hasta que la etiqueta lo haya sido explicada ampliamente. (TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

For MEDICAL AND TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-800-331-2967

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
In case of emergency call toll free the Bayer Environmental Science Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.	
Note To Physician: No specific antidote is available. Treat the patient symptomatically.	

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep children and pets off treated area until dry.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If container is leaking or material spilled for any reason or cause, carefully drain up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer Environmental Science Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer Environmental Science Emergency Response Telephone No. is 1-800-334-7577 or contact Chemtrec at 800-424-9300.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill, by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or vapor. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Keep children or pets off treated area until spray is dry.

Personal Protective Equipment (PPE):

WPS USES: Applicators and Other Handlers who handle this product for any use covered by the Worker Protection Standard (40 CFR part 170) – In general, agricultural plant uses e.g., use of sod farms, must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment. PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Non-WPS USES: Applicators and other handlers who handle this product for any use NOT covered by the Worker Protection Standard (40 CFR part 170) – In general, only agricultural plant uses are covered by the WPS, must wear:

- Shirt and pants
- Gloves
- Shoes plus socks

Engineering controls statements:

• When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Do not formulate this product into other end-use products.

[LINK TO LABEL](#)

DYLOX 420 SL



Dylox[®]
420 SL

TURF AND ORNAMENTAL INSECTICIDE

For control of listed insects through applications to landscape turf and to soil at the bases of landscape flowers, shrubs, and trees.
For turf use on golf courses and on recreational and residential turf areas only.
For Control of White Grubs, Mole Crickets, Sod Webworms, Cutworms and Other Listed Pests of Turfgrass. Controls damaging insects above and below the soil surface.

ACTIVE INGREDIENT:
Trichlorfon, Dimethyl (2,2,2-trichloro-1-hydroxy-ethyl)phosphonate.....37.30%
OTHER INGREDIENTS:.....62.70%
TOTAL:.....100.00%
*3.5 lb trichlorfon per gal of formulated product
EPA Reg. No. 432-1464

KEEP OUT OF REACH OF CHILDREN
CAUTION

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-800-331-2867

Specimen Label
432-1464 160509AV1

APPLICATIONS		
SITE	INSECT	APPLICATION RATE
Landscape Ornamentals (including flowers, shrubs, and trees) and Recreational Lawns & Turf	Cutworms Armyworms	Webworms, incl. sod webworm Cranberry girdler
	Annual bluegrass weevil larvae	
	Annual bluegrass weevil (adults) Billbug larvae Mole crickets Chinch bugs White grubs (including larvae of Asiatic garden beetle, black turfgrass atenioides, European chafer, green June beetle, Japanese beetle, <i>Aphodius</i> spp., May or June beetles (<i>Phyllophaga</i> spp.), northern & southern masked chafers, sugarcane grub (<i>Tomarus</i> spp.), and Oriental beetle)	
<p>Mix specified dosage of DYLOX 420 SL Turf and Ornamental Insecticide in sufficient water to provide adequate distribution during application to lawn, turf, or groundcover area. To ensure that the active ingredient reaches the soil surface, apply with a minimum of 2 gal of water per 1,000 sq ft.</p> <p>Minimize the amount of thatch present prior to application. After application (on the same day), irrigate with sufficient water to thoroughly wet the remaining thatch and top layers of the soil. If sufficient rainfall to thoroughly wet the thatch and top layers of soil occurs after application and prior to irrigation, further irrigation is not required. Under dry conditions, pre-water prior to application for optimum control of white grubs and weevil larvae.</p> <p>If applied to newly mowed <i>Poa annua</i> irrigate as soon as possible after application, particularly under conditions of high temperature and humidity. Potential injury may be exacerbated by tank mixing with other products, or by the addition of surfactants or other additives.</p> <p>If the turf is being maintained at a mowing height of greater than 1 inch, then the maximum specified application rate (6.9 fl oz per 1,000 sq ft) may be required during periods of high pest pressure. Do not use DYLOX 420 SL Turf and Ornamental Insecticide in hose-proportioned sprayers. Note: Do not use treated area or clippings from treated areas for feed or forage. For ornamentals, apply to soil around the plant.</p>		
Narcissus	Narcissus bulb fly	37 fl oz
Apply specified dosage in 100 gallons of water as a drench per 1,000 feet of row. Direct stream to base of plants at beginning of adult fly activity (early May to June). Repeat treatments annually.		

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.
Pesticide Storage: Protect from freezing. Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.
Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, begin immediate measures to contain the liquid by use of a spill kit or absorbent clay carrier and carefully sweep material into a pile preventing runoff into drains or waterways.
Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticide as directed below. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer Environmental Science Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer Environmental Science Emergency Response Telephone No. is 1-800-334-7577, or contact Chemtrec at 1-800-424-9300.
Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.
Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.
Do not use container in connection with food, feed, or drinking water. Completely empty container into application equipment.
Offer for recycling, if available, or dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

[LINK TO LABEL](#)

LESCO CROSSCHECK PLUS

10058333

Net Contents: 1 Gallon
#091624



LESCO
CrossCheck
plus
MULTI-INSECTICIDE

To control pests indoors and outdoors on residential, institutional, public, commercial, and industrial buildings, greenhouses, food handling establishments, and lawns, ornamentals, parks, recreational areas and athletic fields.

When used as a termiticide, individuals/firms must be licensed by the state to apply termiticide products. States may have more restrictive requirements regarding qualifications of persons using this product. Consult the pest control regulatory agency of your state prior to use of this product.

Provides up to 1 month residual control of house flies. Kills fleas for up to 3 months.

EPA Reg. No. 279-3206-10404
EPA Est. 279-NY-1

Active Ingredient:	By Wt.
Bifenthrin*	7.9%
Other Ingredients:	92.1%
	100.0%

KEEP OUT OF REACH
OF CHILDREN
CAUTION

See other panels for additional
precautionary information.

Distributed By:
LESCO, Inc.
1385 East 36th Street
Cleveland, Ohio 44114-4114

15084989

LESCO CrossCheck® Plus Multi-Insecticide contains
3/5 pound active ingredient per gallon.
*Cis isomers 97% minimum, trans isomers 3%
maximum.

Job 51426

[LINK TO LABEL](#)

ROUND UP PRO

Always refer to the label on the product before using Monsanto or any other agricultural product.

2113686-26/CG



The complete broad-spectrum postemergence professional herbicide for industrial, turf and ornamental weed control.

Complete Directions for Use

EPA Reg. No. 524-475

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION IS LIKELY TO RESULT.

2003-1

Read the entire label before using this product.

Use only according to label instructions.

Not all products recommended on this label are registered for use in California. Check the registration status of each product in California before using.

Read the "LIMIT OF WARRANTY AND LIABILITY" statement at the end of the label before buying or using. If terms are not acceptable, return at once unopened.

THIS IS AN END-USE PRODUCT. MONSANTO DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. SEE INDIVIDUAL CONTAINER LABEL FOR REPACKAGING LIMITATIONS.

1.0 INGREDIENTS

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt 41.0%

OTHER INGREDIENTS (including surfactant): 59.2%
100.0%

*Contains 480 grams per liter or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 350 grams per liter or 3 pounds per U.S. gallon of the acid, glyphosate.

This product is protected by U.S. Patent Nos. 5,683,958; 5,703,015; 6,062,238; 6,121,199; and 6,152,000. No license granted under any non-U.S. patent(s).

2.0 IMPORTANT PHONE NUMBERS

1. FOR PRODUCT INFORMATION OR ASSISTANCE IN USING THIS PRODUCT, CALL TOLL-FREE:
1-800-332-3111.

2. IN CASE OF AN EMERGENCY INVOLVING THIS PRODUCT OR FOR MEDICAL ASSISTANCE, CALL COLLECT, DAY OR NIGHT,
(314) 694-4000.

3.0 PRECAUTIONARY STATEMENTS

3.1 Hazards to Humans and Domestic Animals

Keep out of reach of children.

CAUTION!

CAUSES EYE IRRITATION.

Avoid contact with eyes or clothing.

FIRST AID: Call a poison control center or doctor for treatment advice.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses if present after the first 5 minutes then continue rinsing eye.

• Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

• You may also contact (314) 694-4000, collect day or night, for emergency medical treatment information.

• This product is identified as Roundup Pro herbicide, EPA Registration No. 524-475.

DOMESTIC ANIMALS: This product is considered to be relatively non-toxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

• Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

• Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

3.2 Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

3.3 Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL CONTAINERS OR SPRAY TANKS). This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product can only be used in accordance with the Directions for Use on this label or in separately published Monsanto Supplemental Labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State

[LINK TO LABEL](#)

14-14-14



Ornamental Long-term 180-Day Fertilizer

14-14-14



GUARANTEED ANALYSIS

Total Nitrogen (N).....	14.0%
8.5% Urea Nitrogen*	
5.5% Ammoniacal Nitrogen	
Available Phosphate (P ₂ O ₅).....	14.0%
Soluble Potash (K ₂ O).....	14.0%
Calcium (Ca).....	5.5%
Magnesium (Mg).....	1.0%
Sulfur (S).....	6.0%
6.0% Combined Sulfur	
Iron (Fe).....	1.0%
0.53% Water Soluble Iron	

Derived from: Polymer-coated urea, Ammonium phosphate, Potassium sulfate, Magnesium sulfate, Iron sucrate, Calcium carbonate

8.5% Slowly available Nitrogen from polymer-coated urea

CAUTION: This product contains Iron which may stain concrete and other surfaces. Product applied to these surfaces should be promptly removed by sweeping or blowing. DO NOT WASH OFF WITH WATER.

Net Wt. 50 lb (22.68 KG)

Product Code: R529403
Lot# C-XXXXXX
Premium Turf Grade: SGN 220

Made in USA
Manufactured by:
Ferti Technologies
155 East Street Wallingford, CT 06492
Rev 1/20

Directions For Use:

Read the entire label before use. Apply only as directed.

Do not apply fertilizer when plants are suffering from stress related to heat, drought or transplanting. Apply only when foliage is dry.

Landscape Beds Annuals, perennial flowers and Roses: Apply as a top-dress at 1.5 lbs/ 100 sq. ft.

Trees and Shrubs: Shrubs, top-dress with 1/2 cup for every foot of drip line diameter. **Trees,** top-dress with 5 cups for each 2 inch DBH (Diameter Breast Height)

Turf: Apply to seed bed or established turf at 7,000 sq. ft. per bag (see state specific restrictions below)

Vegetable gardens: When seeding or planting, apply evenly around the plants or rows at a rate of 1 lb /100 sq. ft. and mix product into soil to a depth of 1 to 4" and water well. After establishment, apply by top-dress or side-dress at 1 lb / 100 sq. ft.

Soil Incorporation: Mix 8lbs / Cubic Yard

Landscape containers: 6" - apply 1/2 tbsp., 8" - apply 1 tbsp., 10" - apply 2 tbsp., 12" - apply 3 tbsp., 17"-apply 8 tbsp., 26"-apply 15 tbsp.

Virginia: This product is being used in accordance with the recommended application rates established by the VA Nutrient Management Standards & Criteria. This lawn fertilizer contains phosphorous and is only for non-agricultural use on (1) turf during its first growing season, (2) on turf areas being repaired or renovated, and (3) on turf where a soil test performed within the last 3 years indicates a phosphorous deficiency. This fertilizer is not for the routine maintenance of turf.

Maryland: This product is being used in accordance with the recommended application rates established by the University of Maryland. This fertilizer contains phosphorous and may not be used on turf in the state of Maryland except when: 1) providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under §8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) reestablishing or repairing a turf area.

In New Hampshire, this turf/lawn fertilizer contains phosphorous and is only for nonagricultural use on newly established turflawns, for reseeding turflawns, for seedling, or for use when a soil test indicate a phosphorous deficiency. Do not exceed 7.0 lbs. of product (<=1 lb of available phosphate) per 1000 sq. ft. annually.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn and sweep any product that lands on the driveway, sidewalk or street, back onto your lawn.



Warning

Causes skin irritation. Causes eye irritation. May cause respiratory irritation.

Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Do not breathe the dust. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in a well-ventilated area. Wear protective gloves/protective clothing/eye protection/face protection. **IF SWALLOWED:** Remove person to fresh air and keep comfortable for breathing. **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. **IF SKIN IRRITATED:** Call a POISON CENTER/doctor if you feel unwell. Rinse immediately. **IF ON SKIN:** Wash with plenty of water. Take off contaminated clothing and wash it before reuse. Get medical advice/attention if you feel unwell. Store locked up. Dispose of contents/container according to local regulations.

Emergency number : CANUTEC: (613) 996-6666 CHEMTREC : 1-800-424-9300

Tick Killz

All Natural TICK KILLZ™
Effective Pest Control for
Ticks & Mosquitoes

Use on Lawns, Shrubs, Beds & Entire Property —
wherever insects dwell

CONTROLS TICKS, MOSQUITOES, FLEAS & OTHER NUISANCE INSECTS ON CONTACT

This ALL NATURAL product is safer to use around children, pets and wildlife when used as directed

**CONCENTRATED STRENGTH FORMULA
Makes 320 GALLONS**

Keep out of reach of children.

64 FL OZ (2 QTS) 1.89 L

All Natural TICK KILLZ™
Effective Pest Control for
Ticks & Mosquitoes

All Natural Tick Killz™ can be used to control ticks and other nuisance insects all year round!

MIXING DIRECTIONS: Add minimum 1 fl. oz. of Tick Killz™ for every 5 gallons of water; up to a maximum of 4 fl. ozs. of Tick Killz™ for 5 gallons of water for areas of heavy infestation. If using a Mist Blower, add minimum 1 fl. oz. of Tick Killz™ for every 1 gallon of water; up to a maximum of 4 fl. ozs. of Tick Killz™ for 1 gallon of water for heavy infestation.

APPLICATION RATES: For best results, apply no less than 1 gallon of Ready-To-Use (RTU) product per 1,000 square feet (sf). For Mist Blower, apply no less than 1/4 gallon RTU product per 1,000 sf.

KILLS AND REPELS: Ticks, Mosquitoes, Fleas, Gnats, Ants, Aphids, Centipedes, Chinch Bugs, Earwigs, Millipedes, Mites, Spiders, Whiteflies and Other Insects. Safe for pollinators when used as directed.

SPRAYING DIRECTIONS: Spray thoroughly the following: lawns, shrubs, perimeter wood lines, and damp, shady areas where insects tend to dwell and breed. Apply every 4-6 weeks for ticks, every 2-3 weeks for mosquitoes and every 4 weeks for all other insects. Product can be re-applied as often as desired. Allow product to dry before re-entering areas where Tick Killz™ has been applied. Natural Repellents, LLC asserts that Tick Killz™ complies with EPA Regulations for 25(b) exempt pesticide ingredients.

CAUTION: Keep out of reach of children and pets. Do not spray on fruit trees or edible crops. Do not spray into bodies of water. Protective eyewear and gloves must be worn when spraying. Avoid skin or eye contact as product may prove to be an irritant. In case of prolonged exposure to skin or eye contact, flush area with water for at least 15 minutes and consult a physician. If swallowed, seek medical attention immediately. **FOR OUTDOOR USE ONLY.** Store in a dry, secure location. Containers can be reused with Tick Killz™ only or triple rinsed, punctured and included with your normal recycling.

Distributed by
Natural Repellents, L.L.C.
P.O. Box 251
Denville, NJ 07834
973-627-4523
Made in USA

Active Ingredients:

Cedarwood Oil	39.7%
Phenyl Ethyl Propionate	14.7%
Peppermint Oil	1.0%
Inert Ingredient: Ethyl Lactate	44.6%
	100%

0121

0 4 0 2 3 2 0 1 3 2 4 9

Deer Pro



DeerPro™ Deer Repellent Concentrate

ACTIVE INGREDIENT:
Putrescent Whole Egg Solids..... 2.6%
OTHER INGREDIENTS:
Other Ingredients..... 97.4%
TOTAL.....100.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID	
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER Have the product container or label with you when calling a poison control center, doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time (NPIC Web site: www.npic.orst.edu).	

EPA Registration No.: 84178-3
EPA Establishment No.: 84178-CT-001

Net Contents: 50 gallons OR 5 gallons OR 2.5 gallons OR 1 gallon (contains 0.2 lbs of putrescent whole eggs/gallon)